

MINIMAL PRIMES OVER PERMANENTAL IDEALS

GEORGE A. KIRKUP

ABSTRACT. In this paper we discuss minimal primes over permanental ideals of generic matrices. We give a complete list of the minimal primes over ideals of 3×3 permanents of a generic matrix, and show that there are monomials in the ideal of maximal permanents of a $d \times 2d - 1$ matrix if the characteristic of the ground field is sufficiently large. We also discuss the Alon-Jaeger-Tarsi Conjecture, using our results and techniques to strengthen the previously known results.

1. INTRODUCTION

1.1. Definitions. Let $A = (a_{i,j})$ be any $n \times n$ matrix over a commutative ring R . The permanent of this matrix is

$$\sum_{\sigma \in S_n} a_{1,\sigma(1)} a_{2,\sigma(2)} \cdots a_{n,\sigma(n)}.$$

This is often described as the formula for the determinant of A without the minus signs.

Fix m, n, d and a field k . We define $I_d(m, n) \subset k[x_{i,j}]$ to be the ideal generated by the permanents of the $d \times d$ submatrices of $M_{m,n} = (x_{i,j})$, a generic $m \times n$ matrix over k . The main goal of this paper is to understand the minimal primes over $I_d(m, n)$.

1.2. Background. Two recent papers discuss this notion. In [5], Laubacher and Swanson carefully analyze $I_2(m, n)$ for all m, n . They use the language of ideals, minimal primes, radical, Gröbner bases, *etc.* to give a very complete understanding of these ideals.

In [7], Yu discusses some relationships between the rank of a matrix and the “perrank” of a matrix, namely the size of the largest submatrix whose permanent is nonzero. He does not use the language of ideals, preferring an entirely set-theoretic approach to begin to answer some combinatorial questions. These were raised in [1] Alon and Tarsi in 1989 and again treated in [2] in 1994.

1.3. Overview. Our techniques will be very algebraic, and will be simple applications of the multilinearity of permanents which is discussed in Section 2. In Section 3 we review relevant results from [5] and give several proofs based on the results of the previous section.

Date: September 16, 2005.

In Section 4 we discuss the main conjecture of Chapter, which is

Conjecture 1. *If $\text{char } k > d$ or $\text{char } k = 0$ and $n \geq 2d - 1$ then the minimal primes containing $I_d(m, n)$ must either contain a column of the generic matrix or the $(d - 1) \times (d - 1)$ permanents of some $m - 1$ rows.*

This would inductively give the minimal primes over the ideals $I_d(m, n)$ for large m, n if we knew the minimal primes over $I_d(m, n)$ for $m, n \leq 2d - 1$. We prove Conjecture 1 for the case $d = 3$ in Section 5. What we are able to show in general is that any prime over $I_d(d, 2d - 1)$ which contains no entry from some row must contain the $(d - 1) \times (d - 1)$ permanents of the other $d - 1$ rows.

We continue in Section 6 where we discuss the ideals $I_d(d, d + 1)$ in detail. Then in Section 7 we focus on $I_3(3, 4)$, paying special attention to 3×4 matrices with no entries vanishing whose maximal permanents do vanish.

In Section 8 we discuss in general the case in which $m, n > d$. We apply these ideas to the case $d = 3$ in Section 9, in which we list the minimal primes over $I_3(4, 4)$. From this we deduce the minimal primes over $I_3(m, n)$ for all m, n .

In Section 10 we relate our conjecture and results to the combinatorial conjectures in [1].

In this section we strengthen the results of [1] and [2].

2. THE MULTILINEARITY OF THE PERMANENT AND THE ALGEBRA OF PERMANENTS

Let $A = (a_{i,j})$ be a $d \times d$ matrix. Then the permanent of M can be expressed as

$$\sum_{i=1}^d x_{i,1} \cdot A_1^{\hat{i}}$$

where $A_1^{\hat{i}}$ is the $(d - 1) \times (d - 1)$ permanent of the submatrix of A obtained by omitting row i and column 1. This is similar to the expansion by minors (Laplace expansion) for determinants. This expansion can obviously be done for any row or column and this multilinearity is the key observation for our entire study of permanent ideals.

We begin with an example of the importance of multilinearity that we will use throughout this Chapter. If A is a $d \times (d - 1)$ matrix, and column vectors $v_1, \dots, v_d$ have the property that $(A v_i)$ has permanent 0 for all i , then the vector $(A^{\hat{1}}, \dots, A^{\hat{d}})$, of the $(d - 1) \times (d - 1)$ permanents of A , is in the kernel of the matrix $(v_1 v_2 \dots v_d)$. Thus, by rank-nullity, either the determinant of $(v_1 v_2 \dots v_d)$ is 0 or the $(d - 1) \times (d - 1)$ permanents of A vanish identically.

Another application concerns the “algebra of permanents.” Fix $d < n$ and consider the ring $S = k[A_{i_1, \dots, i_d}]$ where $1 \leq i_j \leq n$ for each j and the $A_{i_1, \dots, i_d}$ are indeterminants. Let $R = k[x_{r,s}]$ where $1 \leq r \leq d$ and $1 \leq s \leq n$. Then there is a map $\phi: S \rightarrow R$ taking $A_{i_1, \dots, i_d}$ to the permanent of the matrix

$(c_{i_1}, \dots, c_{i_d})$ where c_j is the j^{th} column of the generic $d \times n$ matrix $(x_{r,s})$. The image of the map $S \rightarrow R$ is the algebra of maximal permanents, and is isomorphic to $S/(\ker \phi)$. We can find interesting elements of $\ker \phi$ in the following way.

Let H be a matrix whose columns are indexed by multi-subsets, $\mathcal{J}$, of $\{1, \dots, n\}$ whose cardinality is $d-1$. Then let the column of H indexed by $\mathcal{J}$ be the vector $A_{1,\mathcal{J}}, \dots, A_{d,\mathcal{J}}$. For example, if $d=3$, and $n=4$ then

$$H = \begin{pmatrix} A_{1,1,1} & A_{1,1,2} & A_{1,1,3} & A_{1,1,4} & A_{1,2,2} & A_{1,2,3} & A_{1,2,4} & A_{1,3,3} & A_{1,3,4} & A_{1,4,4} \\ A_{1,1,2} & A_{1,2,2} & A_{1,2,3} & A_{1,2,4} & A_{2,2,2} & A_{2,2,3} & A_{2,2,4} & A_{2,3,3} & A_{2,3,4} & A_{2,4,4} \\ A_{1,1,3} & A_{1,2,3} & A_{1,3,3} & A_{1,3,4} & A_{2,2,3} & A_{2,3,3} & A_{2,3,4} & A_{3,3,3} & A_{3,3,4} & A_{3,4,4} \\ A_{1,1,4} & A_{1,2,4} & A_{1,3,4} & A_{1,4,4} & A_{2,2,4} & A_{2,3,4} & A_{2,4,4} & A_{3,3,4} & A_{3,4,4} & A_{4,4,4} \end{pmatrix}.$$

We now focus our attention on a particular column, with index $\mathcal{J}$. Then let $A_{\mathcal{J}}^i$ be the permanent of the $(d-1) \times (d-1)$ matrix with columns $\mathcal{J}$ and omitting row i . Then the $\mathcal{J}^{\text{th}}$ column can be expanded as

$$\begin{pmatrix} \sum_i x_{i,1} A_{\mathcal{J}}^i \\ \sum_i x_{i,2} A_{\mathcal{J}}^i \\ \vdots \\ \sum_i x_{i,n} A_{\mathcal{J}}^i \end{pmatrix}.$$

Therefore, no matter what $\mathcal{J}$ is, the column above is in the span of

$$\begin{pmatrix} x_{1,1} \\ x_{1,2} \\ \vdots \\ x_{1,n} \end{pmatrix}, \dots, \begin{pmatrix} x_{d,1} \\ x_{d,2} \\ \vdots \\ x_{d,n} \end{pmatrix}$$

so the rank of H is at most d . Therefore, the $(d+1) \times (d+1)$ minors of H vanish.

3. $I_2(2,3)$

This section reviews a result of Laubenbacher and Swanson [5], and gives three different proofs of it, suggesting three different approaches to the problem in general.

Theorem 2. *If $\text{char } k \neq 2$ then any prime containing $I_2(2,3)$ either contains an entire row of $M_{2,3}$ or it contains an entire column of $M_{2,3}$.*

Proof. Let $\mathfrak{p}$ be a prime over $I_2(2,3)$. The first argument we give is closely related to the computational proof in [5]. If there is no entry of the first row of $M_{2,3}$ in $\mathfrak{p}$ then since $I_{2,3}$ is homogeneous in each row and each column, we can dehomogenize each column so that the $x_{1,1} = x_{1,2} = x_{1,3} = 1$. Then $I_2(2,3)$ becomes $\langle x_{2,1} + x_{2,2}, x_{2,1} + x_{2,3}, x_{2,2} + x_{3,2} \rangle$. Since $\text{char } k \neq 2$, this ideal is generated by $\langle x_{2,1}, x_{2,2}, x_{2,3} \rangle$. Therefore, there must be some entry of $M_{2,3}$ in $\mathfrak{p}$. We may reindex so that $x_{1,1} \in \mathfrak{p}$. Thus $\mathfrak{p}$ must also contain

$x_{2,1}x_{1,2}$ and $x_{2,1}x_{1,3}$ so either $\mathfrak{p}$ contains the first row or the first column of $M_{2,3}$.

The next two proofs use our notation from the previous section,

$$A_{i,j} = x_{1,i}x_{2,j} + x_{1,j}x_{2,i}.$$

Since $A_{1,2} = A_{1,3} = 0$ modulo $I_2(2,3)$, the vector $(x_{1,1}, x_{2,1})$ is in the kernel of the matrix of 1×1 permanents

$$T_{2,3} = \begin{pmatrix} x_{1,2} & x_{1,3} \\ x_{2,2} & x_{2,3} \end{pmatrix}.$$

Therefore, $\langle x_{1,1}, x_{2,1} \rangle \cdot \langle \det T_{2,3} \rangle \in I_2(2,3)$ so $\mathfrak{p}$ either contains a column of $M_{2,3}$ or the 2×2 minors of $M_{2,3}$. Since $\text{char } k \neq 2$, we can combine the minors with the permanents to form all polynomials of the form $x_{1,i}x_{2,j}$ with $i \neq j$. The result then follows similarly to the end of the first proof.

The last proof uses the fact that the symmetric matrix

$$H_{2,3} = \begin{pmatrix} A_{1,1} & A_{1,2} & A_{1,3} \\ A_{1,2} & A_{2,2} & A_{2,3} \\ A_{1,3} & A_{2,3} & A_{3,3} \end{pmatrix}$$

has rank 2 by the discussion in Section 2. Since $A_{i,j} = 0$ modulo $I_2(2,3)$ whenever $i \neq j$, the determinant of $H_{2,3}$ is $A_{1,1} \cdot A_{2,2} \cdot A_{3,3}$. Therefore, one of the $A_{i,i}$ must be in $\mathfrak{p}$. Since $\text{char } k \neq 2$, $A_{i,i} \in \mathfrak{p}$ implies that $x_{1,i} \cdot x_{2,i} \in \mathfrak{p}$ and thus one of these entries is in $\mathfrak{p}$. The proof again follows in the same manner as the first proof. $\square$

These three proofs of the result reflect three different perspectives on how to proceed with the study of permanent ideals.

4. MONOMIALS IN $I_d(d, 2d - 1)$

In this section we will focus on $d \times (2d - 1)$ matrices whose maximal permanents vanish, showing that there must be some entries which vanish (under mild hypotheses) and giving some general tools for understanding the minimal primes over the ideal $I_d(d, 2d - 1)$. We use the approach in the first proof of Theorem 2. The case of $d \times n$ matrices with $n < 2d - 1$ is less understood.

The discussion in this section will revolve around the following conjecture, which will be proved in the case $d = 3$ in Section 5

Conjecture 3. *Let k be any field with $\text{char } k > d$ or 0 and $M_{d,2d-1}$ be the generic $d \times (2d - 1)$ matrix over k . Any minimal prime over $I_d(d, 2d - 1)$ either contains a column of $M_{d,2d-1}$ or contains the $(d - 1) \times (d - 1)$ permanents of some set of $d - 1$ rows of $M_{d,2d-1}$. If $\text{char } k \leq d$ then, in addition to these possibilities, a minimal prime may instead contain the $d \times d$ minors of $M_{d,2d-1}$.*

First we set up some notation for the section. Let $M_{d,n} = (x_{i,j})$ be the generic $d \times n$ matrix where $n > d$. If

$$\overline{\alpha} = (\alpha_1, \dots, \alpha_{d-1})$$

is a strictly increasing sequence with $1 \leq \alpha_1$ and $\alpha_{d-1} \leq 2d-1$ then let $A_{\bar{\alpha}}$ be the $(d-1) \times (d-1)$ subpermanent of M obtained by omitting the first row and including columns $\alpha_1, \dots, \alpha_{d-1}$. Now define

$$A'_{\bar{\alpha}} = A_{\bar{\alpha}} \cdot \prod_{i \notin \bar{\alpha}} x_{1,i}.$$

If

$$\bar{\beta} = (\beta_1, \dots, \beta_d)$$

is a strictly increasing sequence with $1 \leq \beta_1$ and $\beta_d \leq n$ then let $P_{\bar{\beta}}$ be the $d \times d$ subpermanent of M obtained by including columns $\beta_1, \dots, \beta_d$, and as above, let

$$P'_{\bar{\beta}} = P_{\bar{\beta}} \cdot \prod_{i \notin \bar{\beta}} x_{1,i}.$$

Note that

$$P'_{\bar{\beta}} = \sum_{\bar{\alpha} \subset \bar{\beta}} A'_{\bar{\alpha}}.$$

The next proposition will use sums of permanents heavily, so let

$$S_i(\bar{\alpha}) = \sum_{\bar{\beta}} P'_{\bar{\beta}}$$

where the sum runs over all $\bar{\beta}$ such that the cardinality of $\bar{\beta} \cap \bar{\alpha}$ is i . Finally, let

$$T_i(\bar{\alpha}) = \sum_{\bar{\alpha}' \cap \bar{\alpha} \text{ is } i} A'_{\bar{\alpha}'}$$

where the sum runs over all $\bar{\beta}$ such that the cardinality of $\bar{\alpha}' \cap \bar{\alpha}$ is i .

Proposition 4. *For any d, n with $2d-1 > n > d$ and any $\bar{\alpha}$ of size $d-1$, the ideal $I_d(d, n)$ contains*

$$(n-d+1)!(A'_{\bar{\alpha}} + (-1)^{n-d} T_{2d-n-2}(\bar{\alpha})).$$

Proof. In this proof we will treat $A'_{\bar{\alpha}}$ as atomic. We will prove the result by constructing an element of $I_d(d, n)$ and checking the number of times each $A'_{\bar{\alpha}}$ appears.

Fix $\bar{\alpha}$ and let $S_i = S_i(\bar{\alpha})$. Let

$$(1) \quad B = \sum_{i=2d-n-1}^{d-1} (-1)^i (i+n-2d+1)! (d-i-1)! S_i.$$

We claim that $B = \pm(n-d+1)!(A'_{\bar{\alpha}} + T_{2d-n-2}(\bar{\alpha}))$.

Each S_i is a sum of various $A'_{\bar{\alpha}'}$. In fact, if the cardinality of $\bar{\alpha}' \cap \bar{\alpha}$ is i , then S_i contains $n - (d-1) - (d-i-1) = n - 2d + i + 2$ copies of $A'_{\bar{\alpha}'}$. On the other hand, S_{i+1} also contains $d-1-i$ copies of $A'_{\bar{\alpha}'}$.

Given any $A'_{\bar{\alpha}'}$, for which the cardinality of $\bar{\alpha}' \cap \bar{\alpha}$ is i , it can only appear in the definition of S_i or S_{i+1} . Thus we can measure its contribution to B from the definition in (1). It appears

$$(-1)^{i+1} (n-2d+i+2)! (d-i-2)! (d-1-i) + (-1)^i (n-2d+i+1)! (d-1-i)! (n-2d+i+2) = 0$$

times. Therefore, the only contribution to B from $A'_{\bar{\alpha}'}$ comes from the $\bar{\alpha}'$ such that the cardinality of $\bar{\alpha} \cap \bar{\alpha}'$ is $2d - n - 2$ or $\alpha' = \alpha$. In the former case, $A'_{\bar{\alpha}'}$ appears in S_{2d-n-1}

$$d - 1 - (2d - n - 2) = n - d + 1$$

times. Therefore, the total number of times $A'_{\bar{\alpha}'}$ appears in the definition of B is

$$\begin{aligned} (-1)^{2d-n-1} (2d - n - 1 + n - 2d + 1)! \cdot (d - (2d - n - 1) - 1)! \cdot (n - d + 1) \\ = (-1)^{n+1} (0)! \cdot (n - d)! \cdot (n - d + 1) \\ = (-1)^{n+1} (n - d + 1)!. \end{aligned}$$

$A'_{\bar{\alpha}}$ appears in S_{d-1} $n - (d - 1)$ times and in no other S_i . Therefore, the number of times $A'_{\bar{\alpha}}$ appears in the definition of B is

$$\begin{aligned} (-1)^{d-1} (d - 1 + n - 2d + 1)! \cdot (d - (d - 1) - 1)! \cdot (n - (d - 1)) \\ = (-1)^{d-1} (n - d)! \cdot (0)! \cdot (n - d + 1) \\ = (-1)^{d-1} (n - d + 1)!. \end{aligned}$$

This completes the proof since B is a sum of polynomials in $I_d(d, n)$ $\square$

The following proposition is proved in exactly the same manner, and shows why the ideal $I_d(d, 2d - 1)$ is of special importance.

Proposition 5. $d! \cdot A'_{\bar{\alpha}} \in I_d(d, 2d - 1)$ for any $d, \bar{\alpha}$.

Proof. We can use the same proof as in Proposition 4. Let

$$(2) \quad B = \sum_{i=0}^{d-1} (-1)^i i! (d - i - 1)! S_i.$$

We claim that $B = d! \cdot A'_{\bar{\alpha}}$.

Given any $A'_{\bar{\alpha}'}$, with the cardinality of $\bar{\alpha}' \cap \bar{\alpha}$ i , it only appears in the definition of S_i and S_{i+1} . Thus we can count the number of times it appears in B from the definition (2); it appears

$$(-1)^{i+1} (i + 1)! (d - i - 2)! (d - 1 - i) + (-1)^i i! (d - 1 - i)! (i + 1) = 0$$

times. Therefore, the only contribution to B from an $A'_{\bar{\alpha}'}$ comes from $A'_{\bar{\alpha}}$, since there is no $i = d$. Thus $B = d! \cdot A'_{\bar{\alpha}}$ as claimed. $\square$

Notice that this generalizes the first proof of Theorem 2. We get the following corollaries.

Corollary 6. If $\text{char } k > d$ then $\prod x_{i,j} \in I_d(d, 2d - 1)$.

Proof. We induct on d . Certainly $x_{1,1} \in I_1(1, 1)$.

Now assume that $\prod x_{i,j} \in I_{d-1}(d - 1, 2d - 3)$. Then it is clear that $\prod x_{i,j} \in I_{d-1}(d - 1, 2d - 1)$. However, the proposition implies that for any $f \in I_{d-1}(d - 1, 2d - 1)$,

$$f \cdot \prod x_{d,j} \in I_{d, 2d-1},$$

proving the result. $\square$

The following corollary states Proposition 5 set-theoretically.

Corollary 7. *Suppose that $\text{char } k > d$ and M is a $d \times (2d - 1)$ matrix over k whose maximal subpermanents vanish. Then for any row, either there is an entry in that row which vanishes, or the $(d - 1) \times (d - 1)$ subpermanents vanish in the other $d - 1$ rows.*

5. THE IDEAL $I_3(3, 5)$

In this section, we will use the results of the previous section to prove Conjecture 3 in the case $d = 3$. If $\text{char } k = 2$, $I_3(3, 5)$ is just the determinantal ideal, which is prime. Therefore, for the rest of this section we assume that $\text{char } k \neq 2$.

For the sake of clarity, we review the results from the previous section in the case $d = 3$. Let $A_{i,j}$ be the permanent of the 2×2 submatrix of $M_{3,5}$, with columns i, j and rows 2, 3. Then Proposition 5 implies that if $\text{char } k = 0$ or $\text{char } k > 3$ then

$$A_{1,2}x_{1,3}x_{1,4}x_{1,5} \in I_3(3, 5).$$

Moreover, Proposition 4 implies that for all k (recall $\text{char } k \neq 2$),

$$A_{1,2}x_{1,3}x_{1,4} - A_{3,4}x_{1,1}x_{1,2} \in I_3(3, 5).$$

All polynomials of these form which can be obtained by permuting the rows and columns of M are also in $I_3(3, 5)$.

Theorem 8. *If $d = 3$ and $\text{char } k \neq 2, 3$ then any minimal prime containing $I_3(3, 5)$ either contains a column of M or the 2×2 permanents of some two rows. If $\text{char } k = 3$ any prime not containing one of these must contain the 2×2 minors of some collection of 4 columns.*

Proof. Let $\mathfrak{p}$ be a prime over $I_3(3, 5)$. For any three columns of M , either $\mathfrak{p}$ contains the determinant of those three columns, or it contains the 2×2 permanents of the other 2 columns by the multilinearity of the permanents. Suppose that there is a 3×3 submatrix whose determinant is not in $\mathfrak{p}$, and reindex so they are columns 3, 4, 5. Then the 2×2 permanents of the first 2 columns are in $\mathfrak{p}$. Therefore, by Theorem 2 either a row or a column of this 3×2 matrix is in $\mathfrak{p}$. If a column is in $\mathfrak{p}$ we are done. If a row is in $\mathfrak{p}$ then we can reindex so $x_{1,1}, x_{1,2} \in \mathfrak{p}$. If $x_{1,i} \in \mathfrak{p}$ for some other i , reindex so it $i = 3$, and then the 2×2 permanents $A_{1,2}, A_{1,3}, A_{2,3}$ (omitting row 1) are in $\mathfrak{p}$ or $x_{1,4}, x_{1,5} \in \mathfrak{p}$. In the latter case, we are done, and in former case, we are also done by Theorem 2. Therefore, we can assume that $x_{1,i} \notin \mathfrak{p}$ for $i = 3, 4, 5$. In this case, by Proposition 4, we again have $A_{1,2}, A_{1,3}, A_{2,3} \in \mathfrak{p}$, which proves the claim.

There remains the case in which the 3×3 minors of $M_{3,5}$ are in $\mathfrak{p}$. In this case, the matrix $H_{3,5}$ from Section 2 also must have rank 2 (modulo $\mathfrak{p}$), so

its 3×3 minors are in $\mathfrak{p}$. Consider three well-selected columns of H ,

$$H' = \begin{pmatrix} A_{1,1,2} & A_{1,1,3} & 0 \\ A_{1,2,2} & 0 & A_{2,2,3} \\ 0 & A_{1,3,3} & A_{2,2,3} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The existence of H' shows that if the 2×2 minors of the last two rows are not in $\mathfrak{p}$ then all the entries of H' are in $\mathfrak{p}$. This implies that 3×3 permanents of the matrix

$$(3) \quad \begin{pmatrix} x_{1,1} & x_{1,2} & x_{1,3} & x_{1,1} & x_{1,2} & x_{1,3} \\ x_{2,1} & x_{2,2} & x_{2,3} & x_{2,1} & x_{2,2} & x_{2,3} \\ x_{3,1} & x_{3,2} & x_{3,3} & x_{3,1} & x_{3,2} & x_{3,3} \end{pmatrix}$$

vanish. If $\text{char } k \neq 2, 3$, then $3! \neq 0$, so we may assume that there are at least two entries from the first row in $\mathfrak{p}$ by Proposition 5. Thus, by the previous paragraph, if $\text{char } k \neq 3$ we have proved the result.

If $\text{char } k = 3$, we can still use Proposition 4. Let $A_{i,j}$ be the 2×2 subpermanent of $M_{3,5}$ obtained by omitting the first row and including columns i, j (possibly $i = j$). We are in the situation in which the maximal minors of $M_{3,5}$ are in $\mathfrak{p}$, but the 2×2 minors of the last two rows are not in $\mathfrak{p}$, which implied that maximal permanents of matrix (3) were in $\mathfrak{p}$. If an entry of (3) were in $\mathfrak{p}$ then the previous paragraph would imply the result. Therefore, we can dehomogenize each column (set $x_{1,1} = x_{1,2} = x_{1,3} = 1$) so Proposition 4 implies that $A_{i,j} = A_{k,l}$ for all $i, j, k, l \leq 3$ by applying the proposition to each set of 4 columns of (3). Therefore, $A_{i,j} - A_{i,k} = 0$ for all i, j, k so the permanent of

$$\begin{pmatrix} x_{2,i} & x_{2,j} - x_{2,k} \\ x_{3,i} & x_{3,j} - x_{3,k} \end{pmatrix}$$

vanishes for all choices of i, j, k . Therefore, either $x_{2,j} = x_{2,k}$ for all j, k or the 2×2 minors of the bottom two rows of 3 vanish. Since this is true for all j, k , the 2×2 minors of the bottom two rows vanish in either case. Since $I_3(3, 5)$ is also homogeneous in each row and none of the entries of (3) vanish, we can dehomogenize the bottom two rows so that $x_{2,1} = x_{3,1} = 1$. Therefore we have specialized the matrix (3) to

$$(4) \quad \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & x_2 & x_3 & 1 & x_2 & x_3 \\ 1 & x_2 & x_3 & 1 & x_2 & x_3 \end{pmatrix}.$$

The 3×3 permanents of this matrix yield the equations

$$2(1 + x_2 + x_2) = 2(1 + x_3 + x_3) = 0$$

so $x_2 = x_3 = 1$. Rehomogenizing, this means that the 2×2 minors of matrix (3) are in $\mathfrak{p}$.

Therefore, for any pair of columns, either the 2×2 minors of those two columns are in $\mathfrak{p}$ or the 2×2 minors of the other three columns are in $\mathfrak{p}$.

This implies that either there is either a column vanishes modulo $\mathfrak{p}$ or the 2×2 minors of four columns of $M_{3,5}$ are in $\mathfrak{p}$. This completes the proof. $\square$

6. MINIMAL PRIMES OVER $I_d(d, d+1)$

Corollary 6 implies that if $\text{char } k > d$ and if the maximal permanents of a $d \times (2d-1)$ matrix vanish, then at least one entry of the matrix is 0. The natural question is whether this is sharp. In particular, if the permanents of a $d \times n$ matrix, with $n < 2d-1$ vanish, does there need to be an entry of the matrix which is 0? The following example shows that at least in the case $d \times (d+1)$, there need not be any entries which vanish. These examples were constructed by making many of the entries in the matrices 1 and solving the resulting equations.

$$\begin{pmatrix} 1 & \cdots & 1 & 1 & 2-3d \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & \cdots & 1 & 1 & 2-3d \\ 1 & \cdots & 1 & 2-2d & (d-2)(d-1) \\ 1 & \cdots & 1 & d & (2d-1)(d-2) \end{pmatrix}$$

For example, if $d = 3$, this is

$$\begin{pmatrix} 1 & 1 & 1 & -7 \\ 1 & 1 & -4 & 2 \\ 1 & 1 & 3 & 5 \end{pmatrix}.$$

Moreover, the following 4×6 matrix (where $i^2 = -1$) has its maximal permanents vanish,

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & -1 & -1 & -1 \\ 1 & 1 & 1 & i & i & i \\ 1 & 1 & 1 & -i & -i & -i \end{pmatrix}.$$

At this point I am not sure whether there are any 4×6 matrices over k with vanishing maximal permanents and no vanishing entries if k does not have a square root of -1 .

We now discuss $I_d(d, d+1)$, the smallest example of an ideals defined by $d \times d$ permanents, in detail. The next section will explicitly go through the calculations in Section 2 in this case.

Let $M = (x_{i,j})$ be a $d \times (d+1)$ generic matrix and let P_j be the permanent of the square matrix obtained by omitting the j^{th} column of M . Let

$$B_{r,p,q} = \frac{\partial}{\partial x_{r,p}} P_q.$$

Notice that $B_{r,p,q} = B_{r,q,p}$ and $B_{r,p,p} = 0$. Moreover, for $p \neq q$, $B_{r,p,q}$ is the $d-1 \times d-1$ subpermanent of M obtained by omitting row r and columns p, q .

Now consider the matrix of partial derivatives

$$L'_p = \begin{pmatrix} B_{1,1,p} & \cdots & B_{1,d+1,p} \\ \vdots & & \vdots \\ B_{d,d+1,p} & \cdots & B_{d,d+1,p} \end{pmatrix}$$

and let L_p be the $d \times d$ matrix obtained by omitting the p^{th} column of L'_p , which is 0. We define f_p to be the determinant of L_p .

Another natural construction for a matrix of partial derivatives holds a row r constant:

$$W_r = \begin{pmatrix} B_{r,1,1} & \cdots & B_{r,1,d+1} \\ \vdots & & \vdots \\ B_{r,d+1,1} & \cdots & B_{r,d+1,d+1} \end{pmatrix}.$$

Let g_r be the determinant of the symmetric matrix W_r .

Notice the similarity between W_r and $H_{2,4}$ from Section 2. However, there is an important difference. Namely, $A_{i,j}$ in $H_{2,4}$ is the permanent of the 2×2 matrix which has columns i, j whereas $B_{r,i,j}$ omits columns i, j .

Proposition 9. *For any i, j , we have $x_{i,j}f_j, x_{i,j}g_i \in I_d(d, d+1)$.*

Proof. By symmetry we may assume that $i = j = 1$.

For $x_{1,1}f_1$, note that for all $j \neq 1$,

$$P_j = \sum_i x_{i,1} B_{i,1,j}.$$

Let e_j be the determinant of the $d-1 \times d-1$ submatrix of L_1 , omitting the first column and the j^{th} row. Now consider the polynomial

$$\begin{aligned} f'_j &= \sum_j (-1)^j e_j P_j \\ &= \sum_j (-1)^j e_j \sum_i x_{i,1} B_{i,1,j} \\ &= \sum_i x_{i,1} \sum_j (-1)^j B_{i,1,j} e_j. \end{aligned}$$

For $i = 1$, the interior sum is just the expansion by minors of $\det L_1 = f_1$. However, for $i \neq 1$, the interior sum is the expansion by minors of the determinant of L_1 where the first column is replaced by the i^{th} . Thus, for $i \neq 1$, the interior sum is 0. Thus $f'_j = x_{1,1}f_1$, so $x_{1,1}f_1 \in I_d(d, d+1)$.

The second statement is proved in exactly the same method, noting that

$$P_j = \sum_p x_{1,p} B_{1,p,j}.$$

(Recall that if $j = p$ then $B_{1,p,j} = 0$.) From here, the proof proceeds exactly as the previous part. $\square$

The previous proposition allows us to make some statements about the minimal primes over $I_d(d, d+1)$. Let

$$J_d = I_d(d, d+1) + \langle f_j, g_i \rangle \subset k[x_{r,s}].$$

Corollary 10. *If Q is any minimal prime containing $I_d(d, d+1)$, then either Q contains a row, a column, or contains J_d .*

Proof. This is immediate from the previous proposition. If Q does not contain some f_j , then it contains $x_{i,j}$ for all i . Thus Q contains the prime ideal generated by $x_{i,j}$ for all i and the $d \times d$ permanent P_j .

On the other hand, if Q does not contain g_i for some i , then Q contains the ideal generated by $x_{i,j}$ for all j . So Q contains a row. $\square$

7. J_3 , A PRIME CONTAINING $I_3(3, 4)$

In this section, we assume that the characteristic of k is 0, although I believe that the results hold (except for the rational point discussion) for all odd characteristics.

For ease of notation, let

$$M = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 \\ y_1 & y_2 & y_3 & y_4 \\ z_1 & z_2 & z_3 & z_4 \end{pmatrix}$$

and let $I_3(3, 4)$ be the ideal of 3×3 subpermanents of M .

Proposition 11. *Let $k = \mathbb{Q}$. The ideal J_3 , defined in Section 6 is a prime ideal of codimension 4, and $I_3(3, 4)$ is the intersection of J_3 , the primes containing a row of M , and those containing a column of M . Therefore, $I_3(3, 4)$ is a radical complete intersection.*

The proofs are due to Singular [4] and Macaulay 2 [3]. Singular will compute the primary decomposition of $I_3(3, 4)$ (using the GTZ algorithm – the SY algorithm did not terminate) in several hours.

The most interesting ideal in $\text{Ass}(I_3(3, 4))$ is certainly J_3 . Since $I_3(3, 4)$ is a complete intersection, we can compute its degree as

$$\deg(J_3) = \deg(I_3(3, 4)) - 3 \deg(P_1) - 4 \deg(P_3) = 81 - 3(1) - 4(3) = 66$$

where P_1 is a prime containing a row of M and P_3 is a prime containing a column of M , which can be verified directly by Macaulay 2 [3]. The singular locus of $I_3(3, 4)$ is contained in the variety defined by J_3 .

In the previous section we exhibited a rational point on the variety defined by J_3 . The question of finding all rational points with all entries non-zero (full support) can be simplified greatly by the observation that multiplication of any row or column by a scalar results in the multiplication of the permanent by that scalar. This is a result of the homogeneity of $I_d(m, n)$.

In particular, we may normalize five entries to obtain a matrix of the form

$$M = \begin{pmatrix} 1 & 1 & 1 & 1 \\ u & a & b & c \\ u & d & e & f \end{pmatrix}.$$

This is a Zariski open set in the variety defined by J_3 , so there are few matrices which cannot be written in this form, and most of those are contained in other irreducible components of $I_3(3, 4)$. Let J' be the ideal generated by the 3×3 subpermanents of M , which is still homogeneous. The variety defined by J' is certainly Zariski dense in that defined by J_3 since it contains the open set of all matrices with full support.

One interesting situation arises when one of the entries, say $a = -u$. This is evidently a special case since $a = -u$ implies that the 2×2 subpermanent in the top left corner is 0. It turns out that the ideal $I_3(M) + \langle a + u \rangle$ has four primary components, and the irreducible varieties they correspond to may be represented in the following ways.

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & b & c \\ 0 & 0 & eb & -ce \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & d & e & f \end{pmatrix},$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ u & -u & 0 & 0 \\ u & u & e & -e \end{pmatrix}$$

and

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ u & -u & u & u \\ u & 0 & e & -u - e \end{pmatrix}$$

Notice that in no case can the matrix have full support.

Thus, we can change coordinates, $A = a + u, B = b + u, \dots F = f + u$ and still invert the variables. The four 3×3 subpermanents of M can be thought of as relations between the 2×2 subpermanents of the matrix consisting of the last two rows of M . In particular, after some reorganizing, we can rewrite the generators of J' as

$$J' = \langle AE + BD - 2u^2, AF + CD - 2u^2, BF + CE - 2u^2, u \cdot (A + B + C + D + E + F - 6u) \rangle.$$

Since we can invert A, B, C , we get the relations

$$E = \frac{2u^2 - BD}{A} \quad F = \frac{2 - CD}{A}$$

and thus

$$D = \frac{u^2 \cdot (-A + B + C)}{BC}.$$

By symmetry, this means that given the u, A, B, C coordinates, the rest of the coordinates are determined (by field operations), as long as A, B, C, u are nonzero. Recall, however, that one of the generators of J_3 , g_3 , is a

relation only among the elements of the first two rows of M . In particular, we have the following relation

$$\begin{aligned} h &= u^2ab + u^2ac + u^2bc + ua^2b + ua^2c + uab^2 + uac^2 + ub^2c + ubc^2 + a^2bc + ab^2c + abc^2 \\ &= u^2(ab + ac + bc) + u(a^2b + a^2c + ab^2 + ac^2 + b^2c + bc^2) + (a^2bc + ab^2c + abc^2) \\ &= \sigma_1\sigma_3 - \sigma_4 \end{aligned}$$

where $\sigma_1 = a + b + c + u$, $\sigma_3 = abc + abu + acu + bcu$, and $\sigma_4 = abc$.

The form h defines a quartic surface in $\mathbb{P}^3$ which is symmetric in all four variables and has fourteen double points. Since it is quadratic in each variable, we can view it as a double cover of $\mathbb{P}^2$. In that light, we can find several rational curves on the surface, by adding constraints which make the discriminant vanish. For example, if $abc(a + b + c) = 0$, then the constant term vanishes, when viewed as a quadratic in u .

We can also view the surface as an elliptic surface over $\mathbb{P}^1$, since h is a cubic over the function field of $\mathbb{P}^1$ (pick any 2 variables to distinguish). In this light, if we can find a rational point on that elliptic curve (corresponding to a rational curve), we can multiply it by integers using the group law on the elliptic curve. This will yield infinitely many rational curves on the surface unless the original point was a torsion point.

Notice that if $A = B$ then the constant term is 0, so there is a unique solution for C in terms of A and B . In particular, we note that the following matrix has its 3×3 subpermanents vanish for any choice of a .

$$\begin{pmatrix} 1 & 1 & 1 & a+2 \\ 1 & a & a & -a(2a+1) \\ a+1 & -a & -a & a(1-a) \end{pmatrix}$$

Moreover, for all $a \neq -2, -1, -1/2, 0, 1$, this yields a matrix with full support that has its 3×3 subpermanents vanish.

Another interesting special case is when $A = 1/B$ since then the leading term vanishes. In that case we can write another one-parameter family of matrices in J ,

$$\begin{pmatrix} 1 & 1 & a+1 & 3(a+1) \\ 1 & a & -a & a^2+a+1 \\ a+2 & -a(2a+1) & a(1-a) & (a-1)(2a+1)(a+2) \end{pmatrix}.$$

In this case, we get a matrix of full support unless $a = -2, -1, -1/2, 0, 1$ or a is a root of $x^2 + x + 1$ or $3 = 0$.

Other information about J_3 that seems interesting is derived from the free resolution, which Macaulay 2 calculates as

$$0 \leftarrow S/J_3 \leftarrow S \leftarrow S^{11} \leftarrow S^{34} \leftarrow S^{42} \leftarrow S^{24} \leftarrow S^6 \leftarrow 0$$

with Betti diagram

total	:	1	11	34	42	24	6
0	:	1	.	.	.	.	.
1	:	.	.	.	.	.	.
2	:	.	4	.	.	.	.
3	:	.	.	.	.	.	.
4	:	.	.	6	.	.	.
5	:	.	4	12	12	.	.
6	:	.	.	.	.	.	.
7	:	.	3	16	30	24	6

The Castelnuovo-Mumford regularity of J_3 is 7 and its Hilbert polynomial is

$$\frac{11}{840}z^7 + \frac{1}{15}z^6 + \frac{1}{30}z^5 + \frac{41}{12}z^4 + \frac{653}{120}z^3 + \frac{1171}{60}z^2 - \frac{2647}{210}z + 7.$$

8. SUBMAXIMAL PERMANENTS

In this section we will introduce some notation that will make the discussion in Section 9 clearer.

Fix d, m, n and let v be a weakly decreasing sequence of positive integers of finite length $l(v)$, for example $(3, 2, 2)$. Let $s(v)$ be the sum of all the entries of v and define

$$d'(v) = s(v) - l(v) + 1$$

and $e(v) = d - d'(v)$.

Suppose that $s(v) \leq m, n$ and $d'(v) \leq d$. Then we define a prime of type v as follows. Let M' be any submatrix of M of size $m' \times n'$ where $m' + n' + e = m + n$. Then consider any ordered partition of the rows and columns of M' such that the sizes of the sets in the partition (both for rows and columns) matches the entries of v , except perhaps for extra one element sets in the partition. Let a_i be the permanent of the $v_i \times v_i$ submatrix of M' given by the rows and columns in the sets of size v_i in the partitions. Then let $\mathbf{p}$ be generated by a_i for all i and all the entries of M' not used in any of the a_i . Any prime that can be obtained in this way is called a prime of type v .

Notice that given a sequence v , if $v' = (v, 1)$ then the primes of type v are the same as the primes of type v' (so long as $s(v') \leq m, n$) so we can somewhat simplify the description of the types by assuming that no entry of v is 1, except in the case of the empty sequence, which we notate by (1) .

For some easy examples of this, suppose that $d = 2$. Primes of type (1) are those which are generated by all the elements of M except for one row or one column. Primes of type (2) are those which are generated by some 2×2 subpermanent and all the entries of M outside this 2×2 block. There are no other kinds of primes of type v in the case $d = 2$ since $d'(v)$ can be at most 2.

Proposition 12. *Fix m, n, d , and suppose that v is a weakly decreasing sequence of positive integers of finite length. If $s(v) \leq m, n$ and $d'(v) \leq d$ then primes of type v contain $I_d(m, n)$.*

Proof. Since it is clear that the primes of type v are indeed prime, we only need to show that the variety of such a prime is contained in the algebraic set defined by $I_d(m, n)$. The next lemma, which is a corollary of the expansion by minors of permanents allows us to reduce to the case in which $d'(v) = d$.

Lemma 13. *Suppose that $m_1 \leq m$, $n_1 \leq n$, and $d_1 \leq d$. If*

$$d - d_1 \geq (m + n) - (m_1 + n_1)$$

then $I_{d_1}(m_1, n_1) \supset I_d(m, n)$.

Thus we assume $d'(v) = d$, and we induct on $l(v)$. If $l(v) = 1$, then $d = v_1$ and a prime of type v is generated by a $d \times d$ subpermanent of M and all the other entries of M , which evidently contains $I_d(m, n)$.

Let $v = (v_1 \dots v_l)$, let $\bar{v} = (v_1 \dots v_{l-1})$, and let $\mathfrak{p}$ be any prime of type v . We can rearrange columns and rows so that $\mathfrak{p}$ is generated by the upper-left $v_l \times v_l$ subpermanent, all the other entries from the first v_l rows and columns, and in the lower right $m - v_l \times n - v_l$ submatrix, $\bar{M}$, is a prime of type $\bar{v}$. Let

$$\bar{d} = d'(\bar{v}) = d - v_l - 1.$$

By induction, $\mathfrak{p}$ contains all the $\bar{d} \times \bar{d}$ subpermanents of $\bar{M}$. Now, consider any $d \times d$ submatrix, T , of M . If T contains the first v_l rows of M , then since $\mathfrak{p}$ contains all the $v_l \times v_l$ subpermanents of the submatrix given by the first v_l rows of T , expansion by minors implies that the permanent of T is contained in $\mathfrak{p}$. Now assume that T does not contain the first v_l rows of M . Then it must contain at least $d - v_l + 1$ rows from the rest of M . However, by induction, $\mathfrak{p}$ contains the $\bar{d} \times \bar{d}$ subpermanents of this lower submatrix. Now we can use Lemma 13 to finish the proof. In this case $m_1 \geq d - v_l - 1$, $m = n = n_1 = d$, and $d_1 = \bar{d} = d - v_l - 1$. Therefore, since

$$d - d_1 = d - (d - v_l - 1) \geq (m + n) - (m_1 + n_1)$$

$\mathfrak{p}$ contains the permanent of T . Thus we have shown that $\mathfrak{p}$ contains $I_d(m, n)$. $\square$

9. THE IDEALS $I_3(m, n)$

We have already completed our discussion of the minimal primes over $I_3(3, n)$. In this section we will discuss the minimal primes over $I_3(4, 4)$. The primes from the previous section which are relevant are those of type (1), (2), (3), and (2,2). In addition, we say a prime is of type (3A) if it contains the ideal J_3 (from section 7) for some 3×4 (resp. 4×3) submatrix and all the entries of the remaining rows (resp. columns).

Theorem 14. *If $\text{char } k > 3$ then the minimal primes over $I_3(4, 4)$ are of type (1), (2), (3), (3A), or (2,2). If $\text{char } k = 3$ then the ideal of 2×2 minors of the generic matrix is the only other minimal prime.*

Proof. If $\mathfrak{p}$ is any prime over $I_3(4, 4) = I$ and $\mathfrak{p}$ contains all the entries from some row or column, then $\mathfrak{p}$ obviously contains a prime of type (3) or (3A). If $\mathfrak{p}$ contains three elements from some row or column (say $x_{1,1}, x_{1,2}, x_{1,3}$) then $\mathfrak{p}$ either contains $x_{1,4}$ or the 2×2 permanents of the matrix

$$\begin{pmatrix} x_{2,1} & x_{2,2} & x_{2,3} \\ x_{3,1} & x_{3,2} & x_{3,3} \\ x_{4,1} & x_{4,2} & x_{4,3} \end{pmatrix}$$

and thus contains a prime of type (2) or (1).

If $\mathfrak{p}$ contains two elements from some row or column (say $x_{1,1}, x_{1,2}$), then $\mathfrak{p}$ either contains $x_{1,3}, x_{1,4}$ or the 2×2 permanents of the first two columns. By Theorem 2 this means $\mathfrak{p}$ either contains a column of $M_{4,4}$ or the entries in a 2×2 block of $M_{4,4}$. The former case has been dealt with already. In the latter case, suppose $\mathfrak{p}$ contains $x_{1,1}, x_{1,2}, x_{2,1}, x_{2,2}$. Let the $A_{i,j}$ be the 2×2 permanent of the bottom two rows and columns i, j . We have the following relations from the 3×3 permanents,

$$x_{1,3}A_{1,4} + x_{1,4}A_{1,3}, x_{2,3}A_{1,4} + x_{2,4}A_{1,3}$$

so either $x_{1,3}x_{2,4} - x_{1,4}x_{2,3} \in \mathfrak{p}$ or $A_{1,3}, A_{1,4} \in \mathfrak{p}$. However, since $x_{1,3}x_{2,4} - x_{1,4}x_{2,3} \in \mathfrak{p}$ already, the former case would imply that there were three elements from some row in $\mathfrak{p}$ (since $\text{char } k \neq 2$). In the latter case, we already have $A_{1,2} \in \mathfrak{p}$, and similarly $A_{2,3}, A_{2,4} \in \mathfrak{p}$. Therefore, either $\mathfrak{p}$ contains a prime of type (2,2) or $\mathfrak{p}$ contains three entries from one of the bottom two rows.

If $\mathfrak{p}$ contains only one element from some row (say $x_{1,1}$) then by Proposition 4 the 2×2 permanents of the first 2 columns and last three rows vanish. Therefore, we can use one of the previous cases.

Finally, if $\mathfrak{p}$ contains no entries of M , then we can dehomogenize each column so $x_{1,i} = 1$ for each i . As above let $A_{i,j}$ be the 2×2 permanents in the last two columns. By Proposition 4 we know $A_{1,2} = A_{3,4}$, $A_{1,3} = A_{2,4}$, and $A_{1,4} = A_{2,3}$. If the 2×2 minors of the first two rows do not vanish, then by the multilinearity of permanents, the 3×3 minors of

$$\begin{pmatrix} 0 & A_{3,4} & A_{2,4} & A_{2,3} \\ A_{3,4} & 0 & A_{1,4} & A_{1,3} \\ A_{2,4} & A_{1,4} & 0 & A_{1,2} \\ A_{2,3} & A_{1,3} & A_{1,2} & 0 \end{pmatrix}.$$

The top left 3×3 minor of this matrix is $2A_{1,2}A_{1,3}A_{1,4}$. Therefore, one of those is in $\mathfrak{p}$, say $A_{1,2}$. This implies that $A_{3,4} \in \mathfrak{p}$ as well. Moreover, this means that $A_{1,3} + A_{2,3} = 0$ so substituting in the matrix H from Section 2

the the 3×3 minors of the following matrix vanish.

$$\begin{pmatrix} A_{1,1} & 0 & A_{1,3} & -A_{1,3} \\ 0 & A_{2,2} & -A_{1,3} & A_{1,3} \\ A_{1,3} & -A_{1,3} & A_{3,3} & 0 \\ -A_{1,3} & A_{1,3} & 0 & A_{4,4} \end{pmatrix}$$

Taking the first three columns and rows 1,2,4 the minor is

$$-A_{1,1}A_{1,3}(-A_{1,3}) - (-A_{1,3})A_{2,2}A_{1,3} = A_{1,3}^2(A_{1,1} + A_{2,2}).$$

Therefore, either $A_{1,3} = 0$ which would mean that the 2×2 permanents of the bottom two rows vanish, and hence an entry vanishes, or $A_{i,i} = 0$ for all i which also implies that entries of the matrix vanish. Since no entries of the matrix vanish, this is a contradiction. Therefore, the 2×2 minors of every pair of rows vanishes, which is to say that the 2×2 minors of M are in $\mathfrak{p}$. If $\text{char } k \neq 3$, this implies that many entries of M vanish, so the result is proved. $\square$

Corollary 15. *If $m, n \geq 4$ and $\text{char } k \neq 2, 3$ then all minimal primes over $I_3(m, n)$ are of type (1), (2), (3), (3A), or (2,2). If $\text{char } k = 3$ then another minimal prime is the ideal of 2×2 minors of $M_{m,n}$.*

Notice that if $m, n \geq 4$ then all the primes of type 1, 2, 3, 3A, and (2,2) are minimal over $I_3(m, n)$.

10. THE ALON-JAEGER-TARSI CONJECTURE

Now we turn our attention to one of the major motivations for the study of permanental ideals, the Alon-Jaeger-Tarsi Conjecture. We will briefly review the results of [1] and then show how our results and conjectures apply to this conjecture.

The Alon-Jaeger-Tarsi conjecture was stated in [1] as

Conjecture 16. *Let A be a nonsingular $d \times d$ matrix over a finite field k with cardinality $q \geq 4$. There exists a vector v in k^n such that both v and Av have no zero component.*

In [1], Alon and Tarsi show that if A is a matrix over k such that when the matrix A is repeated $q - 2$ times, the resulting matrix $A' = (A \mid A \mid \cdots \mid A)$ has a nonvanishing maximal permanent then there is such a vector. They prove Conjecture 16 for $q = p^k$ where p is prime and $k \geq 2$. In [2] the authors extend this result to the cases $q \geq d + 1 \geq 4$ and $q \geq d + 2 \geq 3$ using a simple counting argument. In Corollary 22 we will show this result for $q \geq d \geq 4$.

Let $A = (a_{i,j})$ be a $d \times n$ matrix over a field k . The key objects of [1] are the polynomials

$$P_A(X_1, \dots, X_n) = \prod_{i=1}^d \left(\sum_{j=1}^n a_{i,j} X_j \right)$$

and

$$P'_A(X_1, \dots, X_n) = X_1 X_2 \cdots X_n \cdot P_A(X_1, \dots, X_d)$$

The definition of P_A immediately implies

Lemma 17. *If $X = (X_1, \dots, X_n)$ is any vector in k^n then $P'_A(X) = 0$ if and only if X and $A(X)$ both have no nonzero entries.*

We introduce the notation of [1] here for convenience and consistency. Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be a vector of nonnegative entries whose sum is d . Then let c_α be the coefficient of the monomial $X_1^{\alpha_1} \cdots X_n^{\alpha_n}$ and let A_α be the $d \times d$ matrix with α_i repetitions of the i^{th} column of A . Claim 1 in [1] is

Lemma 18. *The permanent of A_α is $c_\alpha \cdot \prod (\alpha_i)!$.*

If one could show that for an invertible matrix A , the $d \times 2d$ matrix $A' = (A \mid A)$ has a non-vanishing maximal permanent, then Conjecture 16 would follow. Therefore, Jeff Kahn conjectured that, indeed, A' has a non-vanishing maximal permanent. Yu discusses this conjecture and has some interesting theorems related to it in [7]. The following conjecture is a refinement of Kahn's by De Loera [6]. It should be noted that Conjecture 3 obviously implies this result set-theoretically.

Conjecture 19. *Let $M_{d,d}$ be the generic $d \times d$ matrix over a field k . Then if I is the ideal of $d \times d$ permanents of the $d \times 2d$ matrix $(M_{d,d} \mid M_{d,d})$, $(\det M_{d,d})^d \in I$.*

If k has characteristic 2, the permanent is the determinant so the conjecture is obvious. If the characteristic of the field is 3 then we have the following refinement, the proof of which is related to the proof of the result in [1].

Proposition 20. *If $\text{char } k = 3$ and I_d is the ideal of $d \times d$ permanents of the $d \times 2d$ matrix $(M_{d,d} \mid M_{d,d})$, then*

$$\prod_{i=1}^d D_i(M_i) \subset I_d$$

where M_i is the $i \times d$ submatrix of $M_{d,d}$ given by the first i rows and $D_i(M_i)$ is the ideal of $i \times i$ minors of M_i .

Proof. Let $\{a_1, \dots, a_{d-1}\}$ be any multiset of columns of $M_{d,d} = M$. Then modulo I_d , the permanent of the matrix with columns $\{i, a_1, \dots, a_{d-1}\}$ is 0 for any i because if there is no number repeated three times, then the permanent is a generator of I_d and if there is a number repeated three times, the 3×3 permanents of those three columns vanish since $6 = 0$. Therefore, by expansion by “minors,” the permanent vanishes. This implies that $D_d(M) \cdot P_{d-1}(M \mid M) \in I_d$ where $P_{d-1}(M \mid M)$ is the ideal of $(d-1) \times (d-1)$ subpermanents of $(M \mid M)$. Thus, we can induct, using the case $d = 1$ which is obvious as the base case. $\square$

This refinement is *not* true for arbitrary fields. In fact, a quick calculation using Macaulay 2 [3] shows that if $k = \mathbb{Q}$ then

$$D_3(M_3) \cdot D_2(M_2) \cdot D_1(M_1) \not\subset I_d(M_3 \mid M_3).$$

Computations using Macaulay 2 also suggest that in characteristic 3,

$$\prod_{j=1}^d D_j(M_d) \subset I_d(M_d \mid M_d),$$

which is stronger than Proposition 20.

The next lemma, whose proof is contained in the proof of the Proposition 20, may be of great use in proving Conjecture 19, especially if one could classify the minimal primes over $I_{d-1}(d, d)$.

Lemma 21. *Let $M_{d,d}$ be the generic $d \times d$ matrix over a field k . Then if I is the ideal of $d \times d$ permanents of the $d \times 2d$ matrix $(M_{d,d} \mid M_{d,d})$,*

$$(\det M_{d,d}) \cdot I_{d-1}(d, d) \in I.$$

Using this Lemma and Corollary 15, we prove

Theorem 22. *Conjecture 16 is true for $q \geq d \geq 4$.*

Proof. Let A be an invertible $d \times d$ matrix. By Lemma 18, there is a vector X with only nonzero coordinates such that AX has nonzero coordinates unless the matrix $(A \mid \cdots \mid A)$, with A repeated $q - 2$ times has vanishing maximal permanents. Therefore, we can assume that this is the case. Since A is invertible, Lemma 21 implies that the $d - 1 \times d - 1$ permanents of $(A \mid \cdots \mid A)$, with A repeated $q - 3$ times, vanish. By repeated application of Lemma 21, we see that the $d - q + 3$ sized permanents of A vanish. Since $q \geq d$, we know that the 3×3 permanents of A vanish. However, by Corollary 15, the only invertible $d \times d$ matrices with its 3×3 permanents vanishing are of type (2,2) with $d = 4$. Therefore, we need only consider A of the form

$$\begin{pmatrix} x_1 & x_2 & 0 & 0 \\ y_1 & y_2 & 0 & 0 \\ 0 & 0 & z_1 & z_2 \\ 0 & 0 & w_1 & w_2 \end{pmatrix}$$

with the added constraint that $(A \mid A)$ has its 4×4 permanents vanishing. This is impossible unless one of the rows vanishes, which contradicts the assumption that A is invertible. $\square$

Now we turn our attention away from invertible matrices, and consider arbitrary matrices which have vectors with only nonzero coordinates such that their images have the same property.

Lemma 18 is useful whenever $\alpha_i < p$ for all i . Otherwise, $(\alpha_i)! = 0$ for some i and thus the lemma is trivial for that α . To get past this, we would need to replace the permanent of $A_\alpha = b_{i,j}$ by another object, defined similarly. Fix α and let S_d be the symmetric group on the set

$\{1, \dots, d\}$. Further, let S_α be the quotient group $S_d/(\prod S_{\alpha_i})$ where S_{α_1} acts on the set $\{1, \dots, \alpha_1\}$, S_{α_2} on the set $\{\alpha_1, \dots, \alpha_1 + \alpha_2\}$, etc. Then the product $b_{1,\sigma(1)}b_{2,\sigma(2)} \cdots b_{d,\sigma(d)}$ is well-defined in the sense that for any two representatives of σ in S_d the above products are equal. Using this notation, we get the following lemma.

Lemma 23.

$$c_\alpha = \sum_{\sigma \in S_d/(\prod S_{\alpha_i})} \prod_{i=1}^d b_{i,\sigma(i)}$$

This lemma is what was implicitly used in [1] to prove Lemma 18.

The last piece of [1] that we will use is

Lemma 24. *Let f be any polynomial in $k[x_1, \dots, x_n]$ where k is the finite field with q elements, $k = \mathcal{F}_q$. f is identically 0 over k if and only if f can be reduced to the zero polynomial by the relations $x_i^q = x_i$.*

We are now ready to prove

Theorem 25. *Suppose that A is any $d \times n$ matrix over the finite field $k = \mathcal{F}_q$ and $q > d + 1$. There is a vector $X \in k^n$ such that neither X nor AX has any 0 entries if and only if no row of A is identically 0.*

Proof. It is clear that if a row of A is identically 0 then every vector in the image of A has an entry that is 0. Therefore, we may assume that there is no vectors X such that neither X nor AX has any 0 entries, and we will prove that A has a row which is identically 0.

By Lemma 17 we know that the polynomial P'_A is identically 0 on k^n . Since $q > d + 1$, Lemma 24 implies that $c_\alpha = 0$ for every α .

We proceed by induction on d , since the result is clear for $1 \times n$ matrices. For $\alpha = (d, 0, \dots, 0)$, $c_\alpha = 0$ implies that there is an entry of the first column of A which is 0.

We will prove that either a row of A is identically 0 or the first column is identically 0. Assume that there are exactly $d > e > 1$ entries of the first column which are 0. Reindex so that $a_{1,1} = \dots = a_{e,1} = 0$. Then let

$$L = \{\alpha \mid \alpha_1 = d - e\}.$$

By Lemma 23, for every $\alpha \in L$,

$$c_\alpha = a_{e+1,1} \cdots a_{d,1} \cdot c'_{(\alpha_2, \dots, \alpha_n)}.$$

where A' is the $e \times (n - 1)$ submatrix of A , omitting the first column and including the first e rows and $c'_{(\alpha_2, \dots, \alpha_n)}$ is the coefficient of $X_2^{\alpha_2} \cdots X_n^{\alpha_n}$ of $P'_{A'}$. If $c'_{(\alpha_2, \dots, \alpha_n)} = 0$ for all α then by induction A' must have a row that is identically 0, but that implies that A also has such a row.

Therefore, one of $a_{e+1,1} \cdots a_{d,1}$ must be 0, so more than e entries of the first column vanish which contradicts the assumption that exactly e entries vanish. Therefore, the first column must be identically 0, but this argument

implies that every column is identically 0, so A is identically 0. Therefore, in any case, A has a row which is identically 0. $\square$

Now we consider the case where $d + 1$ is a prime, and use Corollary 7 to establish the following theorem.

Proposition 26. *Suppose that A is any $d \times n$ matrix over the finite field $k = \mathcal{F}_{d+1}$ where $d + 1$ is prime. Further, suppose that there is no vector $X \in k^n$ such that neither X nor AX has any 0 entries. Then either a row of A is identically 0 or*

$$A = \begin{pmatrix} 1 & b_1 & 0 & \dots & 0 \\ 1 & b_2 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & b_d & 0 & \dots & 0 \end{pmatrix}$$

up to scaling of the rows and permutation of the columns, where $\{b_1, \dots, b_d\}$ is the multiplicative group k^ .*

Proof. Since P'_A is identically 0, by Lemmas 17 and 18, the matrix $(A \mid \dots \mid A)$ where A is repeated $d - 1$ times has its maximal permanents vanish. By Corollary 7 either all but $2d - 2$ entries in every row of this matrix vanish or the $(d - 1) \times (d - 1)$ permanents of some $d - 1$ rows vanish. In the latter case, a row of A must vanish by Theorem 25. In the former case, at most 2 entries from every row of A can be nonzero. If any entry of a column is zero, either that column is identically 0 or a row of A is identically 0 by the same argument as in Theorem 25. Therefore, we can assume that exactly two columns of A are nonzero, and so we can reindex A and normalize so that the first column is identically 1. Then the coefficient $c_{i,j,0,\dots,0}$ is the j^{th} elementary symmetric function of the entries in the second column. Since these are 0 for every $0 < j < d$ and

$$c_{d,0,\dots,0} + c_{0,d,\dots,0} = 0$$

so $c_{0,d,\dots,0} = -1$, the entries of the second column of A are the roots of the polynomial $z^{p-1} - 1$, which are all the elements of the multiplicative group k^* . $\square$

In fact, this result holds without the assumption that $q + 1$ is a prime. This can be proven by using the proof above to show that after normalizing one column, the other columns must contain all the nonzero elements of the field. Then, assuming that the first column is identically 1, and the next two columns are nonzero, computing c_α for $\alpha = (i, q - i, 1, 0, \dots, 0)$ for each i , one can show via the Vandermonde determinant that either column 2 or 3 must be zero.

11. CONCLUSION

Proving Conjecture 3 it would be a great step forward in understanding permanental ideals and permanents in general. For example, it would make

it relatively easy to decide whether a matrix of size $d \times n$ where $n \geq 2d - 1$ had vanishing maximal permanents.

It is somewhat striking that the only techniques used in this paper rely on the multilinearity of permanents with some help from symmetry. This suggests that the multilinearity techniques could be successfully applied to ideals defined by other multilinear functions on the $d \times d$ submatrices. I believe that this could be a fruitful approach, but have not had time to pursue this direction.

Another direction I have not pursued is the full primary decomposition of the ideals. Proposition 11 (proved using Macaulay 2) says that $I_3(3, 4)$ is radical, so there are no embedded components in that case. For $I_3(3, 5)$ on the other hand, there are over 200 embedded components, and the decomposition is at this point inscrutable.

REFERENCES

- [1] N. Alon and M. Tarsi, *A nowhere-zero point in linear mappings*, *Combinatorica* **9** (1989), no. 4, 393–395.
- [2] R. D. Baker, J. Bonin, F. Lazebnik, and E. Shustin, *On the number of nowhere zero points in linear mappings*, *Combinatorica* **14** (1994), no. 2, 149–157. MR **MR1289069** (95k:11160)
- [3] Daniel R. Grayson and Michael E. Stillman, *Macaulay 2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [4] G.-M. Greuel, G. Pfister, and H. Schönemann, *SINGULAR 2.0*, A Computer Algebra System for Polynomial Computations, Centre for Computer Algebra, University of Kaiserslautern, 2001, <http://www.singular.uni-kl.de>.
- [5] Reinhard Laubenbacher and Irena Swanson, *Permanental ideals*, *J. Symbolic Computation* **30** (2000), 195–295.
- [6] Bernd Sturmfels, *Solving systems of polynomial equations*, CBMS Regional Conference Series in Mathematics, vol. 97, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2002. MR **2003i:13037**
- [7] Yang Yu, *The permanent rank of a matrix*, *J. Combin. Theory Ser. A* **85** (1999), no. 2, 237–242. MR **MR1673948** (99j:15013)

UNIVERSITY OF CALIFORNIA, BERKELEY
E-mail address: `kirkup@math.berkeley.edu`